

ASTHMA ACTION PLAN

Take this ASTHMA ACTION PLAN with you when you visit your doctor

NAME

DATE

NEXT ASTHMA CHECK-UP DUE

DOCTOR'S CONTACT DETAILS

EMERGENCY CONTACT DETAILS

Name

Phone

Relationship



WHEN WELL *Asthma under control (almost no symptoms)*

ALWAYS CARRY YOUR RELIEVER WITH YOU

Your preventer is:

(NAME & STRENGTH)

Take puffs/tablets times every day

☐ Use a spacer with your inhaler

Your reliever is:

(NAME)

Take puffs

When: You have symptoms like wheezing, coughing or shortness of breath

☐ Use a spacer with your inhaler

Peak flow* (if used) above:

OTHER INSTRUCTIONS

(e.g. other medicines, trigger avoidance, what to do before exercise)



WHEN NOT WELL *Asthma getting worse (needing more reliever e.g. more than 3 times per week, waking up with asthma, more symptoms than usual, asthma is interfering with usual activities)*

Keep taking preventer:

(NAME & STRENGTH)

Take puffs/tablets times every day

☐ Use a spacer with your inhaler

Your reliever is:

(NAME)

Take puffs

☐ Use a spacer with your inhaler

Peak flow* (if used) between and

OTHER INSTRUCTIONS

(e.g. other medicines, when to stop taking extra medicines)

☐ Contact your doctor



IF SYMPTOMS GET WORSE *Asthma is severe (needing reliever again within 3 hours, increasing difficulty breathing, waking often at night with asthma symptoms)*

Keep taking preventer:

(NAME & STRENGTH)

Take puffs/tablets times every day

☐ Use a spacer with your inhaler

Your reliever is:

(NAME)

Take puffs

☐ Use a spacer with your inhaler

Peak flow* (if used) between and

OTHER INSTRUCTIONS

(e.g. other medicines, when to stop taking extra medicines)

Prednisolone/prednisone:

Take each morning for days

☒ Contact your doctor today



DANGER SIGNS

Asthma emergency (severe breathing problems, symptoms get worse very quickly, reliever has little or no effect)

**DIAL 000 FOR
AMBULANCE**

Peak flow (if used) below:

Call an ambulance immediately
Say that this is an asthma emergency
Keep taking reliever as often as needed

NationalAsthma
CouncilAustralia
leading the attack against asthma

www.nationalasthma.org.au

* Peak flow not recommended for children under 12 years.

ASTHMA ACTION PLAN

what to look out for

WHEN WELL



THIS MEANS:

- you have no night-time wheezing, coughing or chest tightness
- you only occasionally have wheezing, coughing or chest tightness during the day
- you need reliever medication only occasionally or before exercise
- you can do your usual activities without getting asthma symptoms

WHEN NOT WELL



THIS MEANS ANY ONE OF THESE:

- you have night-time wheezing, coughing or chest tightness
- you have morning asthma symptoms when you wake up
- you need to take your reliever more than usual eg. more than 3 times per week
- your asthma is interfering with your usual activities

IF SYMPTOMS GET WORSE



THIS MEANS:

- you have increasing wheezing, cough, chest tightness or shortness of breath
- you are waking often at night with asthma symptoms
- you need to use your reliever again within 3 hours

THIS IS AN ASTHMA ATTACK

DANGER SIGNS



THIS MEANS:

- your symptoms get worse very quickly
- you have severe shortness of breath, can't speak comfortably or lips look blue
- you get little or no relief from your reliever inhaler

**CALL AN AMBULANCE IMMEDIATELY: DIAL 000
SAY THIS IS AN ASTHMA EMERGENCY.**

**DIAL 000 FOR
AMBULANCE**

ASTHMA MEDICINES

PREVENTERS

Your preventer medicine reduces inflammation, swelling and mucus in the airways of your lungs. Preventers need to be taken **every day**, even when you are well.

Some preventer inhalers contain 2 medicines to help control your asthma (combination inhalers).

RELIEVERS

Your reliever medicine works quickly to make breathing easier by making the airways wider.

Always carry your reliever with you – it is essential for first aid. Do not use your preventer inhaler for quick relief of asthma symptoms unless your doctor has told you to do this.

To order more Asthma Action Plans visit the National Asthma Council website. A range of action plans are available on the website – please use the one that best suits your patient.

www.nationalasthma.org.au

NationalAsthma
CouncilAustralia
leading the attack against asthma